

**This product is a Hospital Disinfectant, bactericidal at *1:43 dilution (3 oz. per gallon of water) according to the AOAC Use-Dilution Test method, modified in the presence of 250 ppm hard water plus 5% organic serum against:

Pseudomonas aeruginosa *Serratia marcescens* *Klebsiella pneumoniae* *Enterobacter aerogenes*
Staphylococcus aureus *Shigella dysenteriae* *Proteus mirabilis* *Escherichia coli*
Salmonella choleraesuis *Streptococcus pyogenes* *Salmonella typhi* *Brevibacterium ammoniogenes*

Use this product on washable hard nonporous surfaces: floors, walls, cabinets, counters (countertops), stoves (stove tops), sinks, appliances, exterior surfaces of microwave ovens and refrigerators, high chairs, tables, non-wooden picnic tables, outdoor (patio) furniture except cushions and wood frames, chairs, desks, telephones, door knobs, garbage cans, bathtubs, shower stalls, shower walls, glass shower doors, tubs and glazed tiles, urinals, toilets, toilet bowl surfaces, chrome fixtures, hard nonporous surfaces made of metal, stainless steel, chrome, glazed porcelain, glazed ceramic, sealed stone, fiberglass and plastic (such as polystyrene, polypropylene) enameled surfaces, finished woodwork, vinyl and plastic upholstery. Test a small area first. Do not use on PVD finished surfaces.

PELIGRO: Si no sabe el inglés, solicite instrucciones a su supervisor antes de usar esta producto.



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PREPARATION OF DISINFECTANT USE-SOLUTION:

**Add 3 oz. per gallon of water (250-ppm hardness).

DISINFECTANT DIRECTIONS: Apply recommended use-solution with a cloth, mop, sponge, or sprayer or by soaking until surfaces are thoroughly wet. Treated surfaces must remain wet for 10 minutes. Wipe dry with a clean cloth or sponge or let air dry. For sprayer applications, use a coarse spray device. Spray with quick full strokes 6-8 inches from surface. Rub with a brush, sponge or cloth. Do not breathe spray. After use, close nozzle. For heavily soiled areas, a preliminary cleaning is required. Prepare a fresh solution daily or more often if use solution becomes diluted or soiled. Rinse all surfaces that come in contact with food such as countertops appliances, tables and stove tops with potable water before reuse. This product should not be used to clean utensils, glassware and dishes.

This product is not for use on medical device surfaces.

***VIRUCIDAL DIRECTIONS:** Follow Disinfection Directions. Treated surfaces must remain wet for 1 minute.

HEAVY DUTY CLEANER DIRECTIONS: Use 8 oz. per gallon of water for heavy duty cleaning.

TO CLEAN AND DEODORIZE: (Use from 1:10 dilution up to full strength.) Apply product to wet the surfaces thoroughly. Let stand for at least one minute, rinse with water or allow to air dry.

TOILET BOWL/URINAL DISINFECTANT AND CLEANER DIRECTIONS: Remove gross filth prior to disinfection.

From concentrate: Add 3-4 oz. to the toilet bowl or urinal and mix. Thoroughly brush or swab over exposed surfaces and under the rim with a cloth, mop, brush or sponge. Allow to stand for 10 minutes and flush.

From use-solution: Empty toilet bowl or urinal and apply 3-4 oz. per gallon use-solution to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer, allow to stand for 10 minutes and flush.

Manufactured For:
CANBERRA CORPORATION
 3610 Holland-Sylvania Rd.
 Toledo, Ohio 43615

MOLD AND MILDEW CONTROL DIRECTIONS: Add 3-4 oz. per gallon of water, to inhibit the growth of mold and mildew and their odors on hard, nonporous surfaces. Thoroughly wet all treated surfaces completely. Let air dry. Repeat application weekly or when growth or odor reappears.

DIRECTIONS FOR SANITIZING NON-FOOD CONTACT SURFACES:

This product is a sanitizer for hard nonporous non-food contact surfaces at 3/4 oz. per gallon of water dilution against *Klebsiella pneumoniae* and *Staphylococcus aureus*. Treated surfaces must remain wet for 60 seconds. Wipe dry with a clean cloth or sponge or allow to air dry.

KILLS HIV-1 ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human immunodeficiency Virus Type 1 (HIV-1) (Associated with AIDS)

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

PERSONAL PROTECTION: Clean-up should always be done wearing protective gloves, gowns, masks and eye protection.

CLEANING PROCEDURE: Blood and other body fluids containing HIV must be thoroughly cleaned from surfaces and objects before application of this product.

CONTACT TIME: Leave surface wet for 1 minute. Use a 10 minute contact time for disinfection against all other bacteria claimed.

DISPOSAL OF INFECTIOUS MATERIALS: Blood, body fluids, cleaning materials and clothing should be autoclaved and disposed of according to local regulations for infectious waste disposal.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER. Corrosive. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin, or on clothing. Wear protective eyewear (goggles, safety glasses or face shield), protective clothing and protective gloves (rubber or chemical resistant) when handling. Do not breathe spray. Harmful or fatal if swallowed. Avoid contamination of food. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

STORAGE AND DISPOSAL

Store in original container in area inaccessible to children. Do not reuse empty container. Securely wrap original container in several layers of newspaper and discard in trash or recycle.



102908



415

FOAMING DISINFECTANT CONCENTRATE

One-Step Acid Cleaner, Disinfectant, Virucide*, Sanitizer, Mildewstat with Organic Soil Tolerance. Plus Disinfectant Hard Water Tolerance at 3 oz. per gallon (250 ppm calculated as CaCo₃) for Hospital, Institutional, Industrial Use. *Virucidal against HIV-1 (AIDS Virus) and Herpes Simplex Types 1 & 2. Effective as a Disinfectant against *Pseudomonas aeruginosa* (*Pseudomonas*), *Staphylococcus aureus* (*Staph*), and *Streptococcus pyogenes* (*Strep*) Removes Hard Water Deposits and Lime Soap Buildups. Heavy Duty Bathroom Cleaner, Mildewstat.

ACTIVE INGREDIENTS	
Octyl decyl dimethyl ammonium chloride.....	0.75%
Dioctyl dimethyl ammonium chloride.....	0.30%
Didecyl dimethyl ammonium chloride.....	0.45%
Alkyl (C ₁₂ , 50%, C ₁₄ , 40%, C ₁₆ , 10%) dimethyl benzyl ammonium chloride.....	1.00%
INERT INGREDIENTS	97.50%
TOTAL	100.00%

EPA REG. NO. 6836-86-8155 EPA EST. NO. 8155-OH-1

Keep Out Of Reach Of Children

DANGER / PELIGRO

See back panel for additional precautionary statements and First Aid.

Net Contents: One Quart (32 Fl. Oz.) 946 ml